arXiv:1208.3755v2 [cond-mat.mes-hall] 15 Oct 2012

Maximum intrinsic spin-Hall conductivity in two-dimensional systems with k-linear spin-orbit interaction

Tsung-Wei Chen[1], *

[1]*Department of Physics, National Sun Yat-sen University, Kaohsiung 80424, Taiwan*

(Dated: October 16, 2012)

We analytically calculate the intrinsic spin-Hall conductivity (ISHC) (σ^z_{xy} and σ^z_{yx}) in a clean, two-dimensional system with generic k-linear spin-orbit interaction. The coefficients of the product of the momentum and spin components form a spin-orbit matrix $\widetilde{\beta}$. We find that the determinant of the spin-orbit matrix det$\widetilde{\beta}$ determines the effective coupling of spin s_z and orbital motion L_z, and thus, explains the physical origin of the sign change of spin-Hall conductivity. This implies that det$\widetilde{\beta}$ exhibits a discriminant for the non-zero ISHC. Importantly, because the determinant of a matrix is independent of a change in coordinates, the effective coupling det$\widetilde{\beta}$ is independent of the direction of the applied electric field. Furthermore, as the ISHC is non-zero, the other discriminant $\Delta\widetilde{\beta}$ is obtained directly from our result. We show that $\Delta\widetilde{\beta}$ is related to the symmetry of energy dispersion. $\Delta\widetilde{\beta}=0$ corresponds to the energy dispersion with constrained 4-fold symmetry, and in this case, ISHC is a universal constant, and $\sigma^z_{xy}=-\sigma^z_{yx}$. When $\Delta\widetilde{\beta}\neq 0$, that is, the energy dispersion lacks constrained 4-fold symmetry, the ISHC in general depends on the spin-orbit interaction, and $\sigma^z_{xy}\neq -\sigma^z_{yx}$. We further show that two maximum values of ISHC can be achieved ($e/8\pi$ and $e/4\pi$) by tuning $\Delta\widetilde{\beta}$. Certain systems that can have $e/4\pi$ are also discussed. By investigating the equilibrium spin current, we find that det$\widetilde{\beta}$ determines the field strength of SU(2) non-Abelian gauge field.

PACS numbers: 71.70.Ej, 72.25.Dc, 73.63.Hs

I. INTRODUCTION

Within condensed matter physics, spintronics has in itself become a strong field for considerable research, owing to not only its potential applications in electronic technologies but also the many fundamental questions that are raised on the physics of electron spin. [1] Particularly, the spin-orbit interaction recently has strongly attracted the attentions of theoreticians and experimenters since it opens up the possibility of manipulating electron (or hole) spin in non-magnetic materials by electrical means. [2, 3] Since the theoretical prediction of the spin-Hall effect, the application of spintronics has seen considerable advancement. It was shown that the Mott-type skew scattering by impurities would result in separation of opposite spin states via the spin-orbit interaction to the impurity atom. [3] This is the extrinsic spin-Hall effect. Nevertheless, it has been found that in p-type [4] (Luttinger model) and n-type [5] (Rashba model) semiconductors, the spin-polarized current (electron or hole) can be generated by the intrinsic spin-orbit interaction in non-magnetic structure in the absence of impurity scattering, which is called the intrinsic spin-Hall effect (ISHC).

The calculation of spin-Hall conductivity (SHC) plays a crucial role in studying the spin-Hall effect as it can be in comparison with experimental result. The extrinsic spin-Hall effect was experimentally discovered in the three-dimensional (3D) n-type GaAs by optical means via spin accumulation at the edges of a sample [6, 7] and in two-dimensional (2D) n-type AlGaAs/GaAs. [8] The experimental value of the magnitude of SHC $[-0.5(\frac{-2/e}{m\Omega})]$ in Ref. [6] is in agreement with its theoretical value $[0.9(\frac{-2/e}{m\Omega})]$. [9] In 2D p-type AlGaAs/GaAs [10], the experimental value of SHC $[2.5(e/8\pi)]$ also agrees with the theoretical value $[1.9(e/8\pi)]$ in order of magnitude. [11]. Particularly, in Ref. [11], the clean limit is considered in the calculation. In 2D n-type InGaN/GaN, the strain-dependent intrinsic spin-Hall effect detected by optical means is explained in terms of SHC in which the strain effect is included. [12] In 3D metal Pt wire, the large ISHC measured electrically throughout the inverse spin-Hall effect at room temperature is $240(\frac{\hbar/e}{\Omega cm})$. [13, 14] It was theoretically explained in Ref. [15] on the basis of the huge Berry curvature [16] near the Fermi level at the L and X symmetry point in the Pt Brillouin zone; the obtained theoretical value of ISHC is $200(\frac{\hbar/e}{\Omega cm})$ in the absence of impurity scattering. Most recently, a large spin-Hall signal is observed at room temperature in FePt/Au multi-terminal devices. [17]

It has been proposed that a strained semiconductor results in k-linear spin-splitting. [18] Nevertheless, strain-induced spin splitting together with the spin-orbit coupling of the host semiconductor can be simplified and expressed in terms of the coefficients of the spin-orbit matrix [see Eq. (4)]. In order to investigate the influences of these complicated spin-orbit interactions on spin-Hall current systematically, in this study, we focus on generic 2D k-linear spin-orbit coupled systems without impurity scattering. We find that the ISHC can be calculated an-

---

*Electronic address: twchen@mail.nsysu.edu.tw

alytically and that its magnitude and sign change can be described in a unified way.

A typical example of a 2D k-linear system is the Rashba-Dresselhaus system. The ISHC has a universal value $e/8\pi$ and it sign changes because of the competition of Rashba and Dresselhaus coupling strengths. [19] Similar to the sign change of ISHC in Rashba-Dresselhaus system, we will show that the ISHC sign in the generic 2D k-linear systems can also be changed by tuning spin-orbit interactions, and that the sign change leads to the fact that ISHC must cross zero. The two results are uniquely described by the determinant of the spin-orbit matrix, $\det\widetilde{\beta}$, defined in the next section [see Eq. (14)]. We also show that $\det\widetilde{\beta}$ physically expressed as the effective coupling of spin z-component s_z and orbital angular momentum L_z. As a result, the vanishing spin-Hall conductivity corresponds to the decoupling of spin and orbital motion.

Importantly, by analytically calculating the ISHC, we find that ISHC can have a value higher than $e/8\pi$ in certain systems. Unlike in the Rashba-Dresselhaus system, we show that the maximum value of ISHC in a generic k-linear system is $e/4\pi$. The variation between $e/8\pi$ and $e/4\pi$ can be obtained by tuning the spin-orbit interaction, such as Rashba coupling controlled effectively by gate voltage [20] and strain-induced splitting by Laser. [12]

Our present paper is organized as follows. In Sec. II, we define the spin-orbit matrix obtained from the coefficients of the product of momentum and spin. The intrinsic spin-Hall conductivity is shown to be proportional to the determinant of spin-orbit matrix $\det\widetilde{\beta}$. We also use Foldy-Wouthuysen transformation to show that the effective coupling of the spin z-component s_z and orbital angular momentum L_z is $-\frac{2m}{\hbar^4}\det\widetilde{\beta}$. In Sec. III, we analytically calculate the ISHC of the generic k-linear system. Our calculation results show that in general the ISHC depends on the strength of spin-orbit couplings, and it is not necessarily a universal constant. Furthermore, we define the other discriminant $\Delta\widetilde{\beta}$ and show that for certain systems, the ISHC can have a maximum value of $e/4\pi$. Realistic systems having higher values of ISHC are also discussed. In Sec. IV, we study the relationship between the symmetry of energy dispersion and the asymmetric properties of ISHC. In Sec. V, we discuss the relationship between equilibrium spin current and spin-orbit matrix. Similar to the ISHC, the equilibrium spin current is related to $\det\widetilde{\beta}$. We show that $\det\widetilde{\beta}$ plays the role of color magnetic field strength. Our conclusions are presented in Sec. VI.

II. INTRINSIC SPIN-HALL CONDUCTIVITY

A. spin-orbit matrix and ISHC

The generic k-linear spin-orbit coupled 2D systems are related to the spin-orbit matrix $\widetilde{\beta}$,

$$\widetilde{\beta} = \begin{pmatrix} \beta_{xx} & \beta_{xy} \\ \beta_{yx} & \beta_{yy} \end{pmatrix}, \tag{1}$$

where the coefficients β_{ij} represent the spin-orbit interactions in 2D systems. As an example, let us consider the pure Bychkov-Rashba system $[\alpha(\sigma_x k_y - \sigma_y k_x)]$ [21], the pure Dresselhaus system $[\beta(\sigma_x k_x - \sigma_y k_y)]$ [22] and the Dirac-type system $[g(\sigma_x k_x + \sigma_y k_y)]$ [23, 24]; the corresponding spin-orbit matrices for these systems are

$$\widetilde{\beta}_R = \begin{pmatrix} 0 & \alpha \\ -\alpha & 0 \end{pmatrix},\ \widetilde{\beta}_S = \begin{pmatrix} \beta & 0 \\ 0 & -\beta \end{pmatrix},\ \widetilde{\beta}_D = \begin{pmatrix} g & 0 \\ 0 & g \end{pmatrix}, \tag{2}$$

respectively. Another example is the spin splitting in a bulk strained semiconductor. [18, 25] The spin-orbit matrices $\widetilde{\beta}_1$ and $\widetilde{\beta}_2$ denote, respectively, the system with structure inversion asymmetry (SIA) strain-induced splitting and the system with bulk inversion asymmetry (BIA) strain-induced splitting. They are given by

$$\begin{aligned} \widetilde{\beta}_1 &= \begin{pmatrix} 0 & \frac{1}{2}C_3\epsilon_{xy} \\ -\frac{1}{2}C_3\epsilon_{yx} & 0 \end{pmatrix}, \\ \widetilde{\beta}_2 &= \begin{pmatrix} D(\epsilon_{zz}-\epsilon_{yy}) & 0 \\ 0 & D(\epsilon_{xx}-\epsilon_{zz}) \end{pmatrix}, \end{aligned} \tag{3}$$

where the structure constant $C_3 > 0$ and $D > 0$. [18, 26] Thus, in addition to SIA and bulk-inversion-symmetry breaking induced spin-orbit interaction, the strain-induced spin splitting is included in the spin-orbit matrix elements. Accordingly, we do not pose any restrictions on the spin-orbit matrix elements in the following calculations. The 2D Hamiltonian can be written as

$$\begin{aligned} &H = \varepsilon_{\mathbf{k}} + H_{so}, \\ &H_{so} = \sum_{ij} \beta_{ij}\sigma_i k_j = (\sigma_x \ \ \sigma_y) \begin{pmatrix} \beta_{xx} & \beta_{xy} \\ \beta_{yx} & \beta_{yy} \end{pmatrix} \begin{pmatrix} k_x \\ k_y \end{pmatrix}, \end{aligned} \tag{4}$$

where $\varepsilon_{\mathbf{k}} = \hbar^2 k^2/2m$ and σ_i $(i = x, y)$ are the Pauli spin matrices. The momenta k_x and k_y are parallel to [100] and [010] directions, respectively. The interplay of various spin-orbit interactions is described by the combination of spin-orbit matrices. For example, the Rashba-Dresselhaus system, the spin-orbit matrix is given by $\widetilde{\beta}_D + \widetilde{\beta}_R$ and the corresponding Hamiltonian is $H = \varepsilon_{\mathbf{k}} + \alpha(\sigma_x k_y - \sigma_y k_x) + \beta(\sigma_x k_x - \sigma_y k_y)$. For calculating ISHC, we further rewrite Eq. (4) in the following form.

$$H = \varepsilon_{\mathbf{k}} + d_x\sigma_x + d_y\sigma_y, \tag{5}$$

where

$$\begin{aligned} d_x &= \beta_{xx}k_x + \beta_{xy}k_y \\ d_y &= \beta_{yx}k_x + \beta_{yy}k_y. \end{aligned} \tag{6}$$

The eigenenergy is $E_{n\mathbf{k}} = \varepsilon_{\mathbf{k}} - nd$ for two branches $n = \pm$, where the dispersion term $d = \sqrt{d_x^2 + d_y^2}$ can be written as

$$d = k\Gamma(\phi), \tag{7}$$

where

$$\Gamma(\phi)^2 = \left(\beta_{xx}^2 + \beta_{yx}^2\right)\cos^2\phi + \left(\beta_{xy}^2 + \beta_{yy}^2\right)\sin^2\phi + \left(\beta_{xx}\beta_{xy} + \beta_{yx}\beta_{yy}\right)\sin(2\phi). \tag{8}$$

The energy dispersion Eq. (8) satisfies $\Gamma(\phi) = \Gamma(\phi + \pi)$ because the time-reversal symmetry is preserved. For a positive chemical potential ($\mu > 0$), the Fermi momenta for two branches satisfy the following condition

$$k_F^+ - k_F^- = \frac{2m\Gamma(\phi)}{\hbar^2}. \tag{9}$$

The ISHC can be evaluated by using the Kubo formula [27]

$$\sigma_{ij}^z(\omega) = \lim_{\omega\to 0}\frac{Q_{ij}^z(\omega)}{-i\omega}. \tag{10}$$

$Q_{ij}^z(\omega)$ is the spin current-charge current correlation function. The index j represents the direction of applied electric field and i the direction of response current. The conventional definition of spin current is $J_i^z = \frac{\hbar}{4}\left\{\partial H/\partial k_i, \sigma_z\right\}$ [28], and charge current is defined as $J_j = e\frac{\partial H}{\partial \hbar k_j}$. From the standard approach, it can be shown that [29]

$$Q_{ij}^z(\omega) = i\omega\frac{e}{V}\sum_{\mathbf{k}}\frac{f_{\mathbf{k}+} - f_{\mathbf{k}-}}{d(\omega^2 - 4d^2)}\frac{\partial\varepsilon_{\mathbf{k}}}{\partial k_i}\left(d_x\frac{\partial d_y}{\partial k_j} - d_y\frac{\partial d_x}{\partial k_j}\right), \tag{11}$$

where $f_{\mathbf{k}\pm}$ represents the Fermi function for two energy branches. Note that the correlation function contains the kinetic term. Next, we focus on spin-Hall conductivity in the static case ($\omega = 0$). When an electric field is applied in y direction ([010]), and the spin-Hall response in x direction ([100]) is given by

$$\sigma_{xy}^z(0) = -\frac{e}{V}\sum_{\mathbf{k}}\frac{f_{\mathbf{k}+} - f_{\mathbf{k}-}}{-4d^3}\frac{\partial\varepsilon_{\mathbf{k}}}{\partial k_x}\left(d_x\frac{\partial d_y}{\partial k_y} - d_y\frac{\partial d_x}{\partial k_y}\right). \tag{12}$$

Substituting Eqs. (6), (7), (8), and (9) into Eq.(12) and using the replacement $\frac{1}{V}\sum_{\mathbf{k}} \to \int\frac{kdkd\phi}{(2\pi)^2}$, after a straightforward calculation, we obtain

$$\sigma_{xy}^z(0) = \frac{e}{8\pi^2}\left(\det\widetilde{\beta}\right)\int_0^{2\pi} d\phi\frac{\cos^2\phi}{\Gamma(\phi)^2}, \tag{13}$$

where $\det\widetilde{\beta}$ stands for the determinant of the spin-orbit matrix $\widetilde{\beta}$

$$\det\widetilde{\beta} = \beta_{xx}\beta_{yy} - \beta_{xy}\beta_{yx}. \tag{14}$$

Equation. (13) indicates that the spin-Hall conductivity vanishes when $\det\widetilde{\beta} = 0$. For example, in the Rashba-Dresselhaus system ($\widetilde{\beta}_R + \widetilde{\beta}_S$), we have $\beta_{xx} = \beta$, $\beta_{xy} = \alpha$, $\beta_{yx} = -\alpha$, $\beta_{yy} = -\beta$, and $\det\widetilde{\beta} = \alpha^2 - \beta^2$. Therefore, when $\alpha^2 = \beta^2$, the spin-Hall conductivity vanishes. It has been shown that the the vanishing spin-Hall conductivity in the Rashba-Dresselhaus system results from the fact that the orbital motion is decoupled from the spin z-component when $\alpha^2 = \beta^2$. [30] We can generalize this argument to a generic k-linear system and show that the decoupling of orbital motion from spin is related to Eq. (14). This can be proved in the following subsection.

B. effective coupling of spin and orbital motion

We apply the Foldy-Wouthuysen transformation [31] to the Hamiltonian Eq. (4), and diagonalize the Hamiltonian up to some order of β_{ij}. Because $\det\widetilde{\beta}$ is order of β_{ij}^2, the unitary transformation that can diagonalize Eq. (4) up to second order is given by (see Appendix A)

$$\begin{aligned} U &= \exp\left\{-i\frac{m}{\hbar^2}D\right\}, \\ D &= \sigma_x F_x + \sigma_y F_y, \\ F_x &= \beta_{xx}x + \beta_{xy}y, \\ F_y &= \beta_{yx}x + \beta_{yy}y, \end{aligned} \tag{15}$$

where F_x and F_y are obtained by using the replacements $k_x \to x$ and $k_y \to y$ in d_x and d_y. It can be shown that

$$[D, \varepsilon_{\mathbf{k}}] = \frac{i\hbar^2}{m}H_{so}. \tag{16}$$

Using the unitary transformation Eq. (15) and Eq. (16), then Eq. (4) becomes (up to the second order of β_{ij})

$$\begin{aligned} H' =& U^\dagger H U \\ =& \varepsilon_{\mathbf{k}} + \frac{im}{\hbar^2}\left(1 - \frac{1}{2!}\right)[D, H_{so}] + o(\beta_{ij}^3). \end{aligned} \tag{17}$$

It can be shown that

$$[D, H_{so}] = i\sum_{ij}\beta_{ij}^2 + \frac{2i}{\hbar}\left(\det\widetilde{\beta}\right)\sigma_z L_z, \tag{18}$$

where $L_z = \hbar(xk_y - yk_x)$ is the orbital angular momentum. Substitute Eq. (18) into Eq. (17), we obtain

$$H' = \varepsilon_{\mathbf{k}} - \frac{m}{2\hbar^2}\sum_{ij}\beta_{ij}^2 + h_D + o(\beta_{ij}^3), \tag{19}$$

where

$$h_D = -\frac{m}{\hbar^3}\left(\det\widetilde{\beta}\right)\sigma_z L_z \tag{20}$$

Equation (20) shows that the coupling between orbital motion L_z and spin z component σ_z is proportional to det$\widetilde{\beta}$. Therefore, Eq. (14) together with Eqs. (13) and (20) exhibits a discriminant for a non-vanishing spin-Hall conductivity:

$$\begin{aligned} \det\widetilde{\beta} = 0 &\rightarrow \sigma^z_{xy}(0) = 0, \\ \det\widetilde{\beta} \neq 0 &\rightarrow \sigma^z_{xy}(0) \neq 0. \end{aligned} \tag{21}$$

Hence, based on the discriminant for k-linear system, we can justify the condition for non-vanishing SHC simply from the spin-orbit matrix $\widetilde{\beta}$ without calculating the SHC.

III. MAXIMUM VALUES OF ISHC

As shown in Sec. II, the determinant of the spin-orbit matrix, det$\widetilde{\beta}$, is the effective coupling of s_z and L_z. Importantly, det$\widetilde{\beta}$ is independent of a change in coordinates, and thus, the strength of the effective coupling is also independent of the direction of the applied electric field. That is, if we change the coordinate (k_x, k_y) to an arbitrary coordinate (k'_x, k'_y), we have $\det\widetilde{\beta} = (\beta_{xx}\beta_{yy} - \beta_{yx}\beta_{xy}) = \det\widetilde{\beta}' = (\beta_{x'x'}\beta_{y'y'} - \beta_{y'x'}\beta_{x'y'})$. However, the direction of the applied electric field may affect the value of ISHC in the sense that the symmetry of energy dispersion is determined by spin-orbit interaction. Therefore, in this section, we evaluate Eq. (13) and study the dependence of spin-orbit interaction on ISHC.

A. discriminant for maximum values

In order to evaluate the integral in Eq. (13), we transform the integral to the contour integral in a complex plane. If z is defined as $z = e^{i\phi}$, the integral becomes a line integral along a closed loop with unit radius. The function $\Gamma(\phi)$ [8] can be rewritten as

$$\Gamma(\phi)^2 \rightarrow \Gamma(z)^2 = \frac{1}{4z^2}\left(\lambda_1 z^2 + \lambda_2\right)\left(\lambda_2^* z^2 + \lambda_1^*\right), \tag{22}$$

where " $*$ " symbolizes the complex conjugate and

$$\begin{aligned} \lambda_1 &= (\beta_{xx} + \beta_{yy}) + i(\beta_{yx} - \beta_{xy}) \\ \lambda_2 &= (\beta_{xx} - \beta_{yy}) + i(\beta_{yx} + \beta_{xy}). \end{aligned} \tag{23}$$

The integral in Eq. (13) can be evaluated by calculating the residue inside the unit circle $|z| = 1$. The conditions for the poles appearing in the unit circle indicate the boundary of change of ISHC in sign. By using the standard residue theorem [32], the result is derived as

$$\int_0^{2\pi} d\phi \frac{\cos^2\phi}{\Gamma(\phi)^2} = \frac{4\pi}{|\lambda_>|^2 - |\lambda_<|^2}\left[1 - Re\left(\frac{\lambda_<}{\lambda_>}\right)\right], \tag{24}$$

where $Re(\cdots)$ represents the real part of a complex number. $\lambda_>$ ($\lambda_<$) is taken from the relative maximum (minimum) value of ($|\lambda_1|$, $|\lambda_2|$). That is, if $|\lambda_1| > |\lambda_2|$ then $\lambda_> = \lambda_1$ and $\lambda_< = \lambda_2$, and vice versa. Equation (24) can be further simplified. Using Eq. (23), we find that

$$|\lambda_1|^2 - |\lambda_2|^2 = 4\det\widetilde{\beta}. \tag{25}$$

If $\det\widetilde{\beta} \neq 0$ in Eq. (25), it cancels det$\widetilde{\beta}$ appearing in Eq. (13). Combining Eq. (13) together with (24) and (25), we have

$$\sigma^z_{xy}(0) = \begin{cases} sgn(\det\widetilde{\beta})\dfrac{e}{8\pi}\left[1 - Re\left(\dfrac{\lambda_<}{\lambda_>}\right)\right] & ; \det\widetilde{\beta} \neq 0 \\ 0 & ; \det\widetilde{\beta} = 0 \end{cases}, \tag{26}$$

where $sgn(\cdots)$ is the sign function, i.e. $sgn(\det\widetilde{\beta}) = \frac{\det\widetilde{\beta}}{|\det\widetilde{\beta}|}$. We have $sgn(\det\widetilde{\beta} > 0) = +1$ and $sgn(\det\widetilde{\beta} < 0) = -1$. The real part of $\lambda_</\lambda_>$ in Eq. (26) can be written in terms of coefficients of spin-orbit matrix,

$$Re\left(\frac{\lambda_<}{\lambda_>}\right) = \frac{(\beta^2_{xx} - \beta^2_{yy}) + (\beta^2_{yx} - \beta^2_{xy})}{\sum_{ij}\beta^2_{ij} + 2|\det\widetilde{\beta}|}. \tag{27}$$

Note that in Eq. (27), there is an absolute value of det$\widetilde{\beta}$. In the case $|\lambda_1| > |\lambda_2|$, we have $\det\widetilde{\beta} > 0$ and $|\det\widetilde{\beta}| = \det\widetilde{\beta}$. In the case $|\lambda_1| < |\lambda_2|$, we have $\det\widetilde{\beta} < 0$ and $|\det\widetilde{\beta}| = -\det\widetilde{\beta}$. We find that $\sum_{ij}\beta^2_{ij}$ can be expressed in term of $\Gamma(\phi)$: $\sum_{ij}\beta^2_{ij} = \frac{1}{2}[\Gamma(\frac{\pi}{4})^2 - \Gamma(-\frac{\pi}{4})^2]$.

Equation (26) is our main result. First, we find that if $\det\widetilde{\beta} \neq 0$, then its sign determines the sign of ISHC. Based on Eq. (25), the change in sign of ISHC can be simply determined by the sign of det$\widetilde{\beta}$, and the physical origin of the sign change is described in Eq. (20). The orbital motion L_z couples to $+\sigma_z$ when $\det\widetilde{\beta} > 0$, while L_z couples to $-\sigma_z$ when $\det\widetilde{\beta} < 0$. In the Rashba-Dresselhaus system, $\lambda_1 = -2i\alpha$, $\lambda_2 = 2\beta$, we have $\det\widetilde{\beta} = (\alpha^2 - \beta^2)$ and the ratio of λ_1 and λ_2 is purely imaginary; the spin-Hall conductivity is a universal constant and changes sign by tuning the relative strength of spin-orbit interactions such that det$\widetilde{\beta}$ changes sign. This result is in agreement with the previous theoretical results [19]. Rashba coupling and Dresselhaus coupling are usually of the same order of magnitude in the GaAs quantum well. [33] In the II-VI semiconductor, Rashba coupling is larger than Dresselhaus coupling, while in the III-V semiconductor, Dresselhaus coupling would be larger than Rashba coupling. [33] In the narrow-gap compounds, Rashba coupling dominates. [34]

In a pure Rashba system, $\det\widetilde{\beta}_R = \alpha^2$ is always positive, while in a pure Dresselhaus system, $\det\widetilde{\beta}_S = -\beta^2$ is always negative. This implies that the same orbital motion couples to an opposite spin z component, and thus, the spin-Hall conductivity of the two systems is different in sign. [35] Unlike the 3D case, where the ISHC depends on Fermi momentum [4], the ISHC in the 2D case is independent of Fermi momentum.

On the other hand, the ISHC generally depends on the spin-orbit interaction [Eq. (27)], and it is not necessarily a universal constant. We note that the denominator of Eq. (27) is always positive. Nevertheless, the numerator of Eq. (27) can be either positive or negative. For convenience in the following discussion, we define

$$\Delta\widetilde{\beta} \equiv (\beta_{xx}^2-\beta_{yy}^2)+(\beta_{yx}^2-\beta_{xy}^2). \tag{28}$$

Equation (28) can also be written as $\Delta\widetilde{\beta} = [\Gamma(0) - \Gamma(\frac{\pi}{2})] = [\Gamma(\pi) - \Gamma(\frac{3\pi}{2})]$. The real part in Eq. (26) is always less than unity in magnitude. This is because $\sum_{ij}\beta_{ij}^2 = \beta_{xx}^2+\beta_{yy}^2+\beta_{xy}^2+\beta_{yx}^2$ is larger than $\Delta\widetilde{\beta}$. For $\Delta\widetilde{\beta} \geq 0$, the ISHC is less than $\frac{e}{8\pi}$. Therefore, the spin-Hall conductivity has an upper bound in magnitude,

$$|\sigma_{xy}^z(0)| \leq \frac{e}{8\pi},\ \Delta\widetilde{\beta} \geq 0. \tag{29}$$

It has been shown that the torque spin-Hall conductivity is opposite in sign and twice as larger in magnitude when compared to the conventional spin-Hall conductivity. [35] The total spin-Hall conductivity only has an overall negative sign. The equality in Eq. (29) is valid only when $Re(\lambda_1/\lambda_2) = 0$ or $Re(\lambda_2/\lambda_1) = 0$; Eq. (27) gives the same result for both conditions: $(\beta_{xx}^2-\beta_{yy}^2)+(\beta_{yx}^2-\beta_{xy}^2) = 0$. In other words, the system satisfies the conditions: (1)$\beta_{xx}^2 = \beta_{yy}^2$ and $\beta_{yx}^2 = \beta_{xy}^2$, (2)$\beta_{xx}^2 = \beta_{xy}^2$ and $\beta_{yx}^2 = \beta_{yy}^2$, or (3) $\beta_{xx}^2+\beta_{yx}^2 = \beta_{yy}^2+\beta_{xy}^2$. Some spin-orbit coupled systems satisfy condition (1), i.e, the pure Rashba, pure Dresselhaus, Rashba-Dresselhaus, and Dirac-Rashba systems.

Interestingly, we find that if $\Delta\widetilde{\beta} < 0$, then $Re\,(\lambda_</\lambda_>)$ is negative and the spin-Hall conductivity satisfies

$$|\sigma_{xy}^z(0)| < \frac{e}{4\pi},\ \Delta\widetilde{\beta} < 0. \tag{30}$$

The spin-Hall conductivity still has an upper bounded value; however, it can exceed the value $e/8\pi$. The three conditions are summarized in Fig. 1, where we define $N = Re(\lambda_</\lambda_>)$ and it has been shown that $|N| < 1$. The spin-Hall conductivity $|\sigma_{xy}^z(0)| = \frac{e}{8\pi}(1-|N|)$ for $\Delta\widetilde{\beta} > 0$, $|\sigma_{xy}^z(0)| = \frac{e}{8\pi}$ for $\Delta\widetilde{\beta} = 0$ and $|\sigma_{xy}^z(0)| = \frac{e}{8\pi}(1+|N|)$ for $\Delta\widetilde{\beta} < 0$.

B. Realistic systems having higher ISHC

We discuss some specific systems that are possible to have $\Delta\widetilde{\beta} < 0$. Consider BIA strain split described by the spin-orbit matrix $\widetilde{\beta}_2$ and assume that $(\epsilon_{zz} - \epsilon_{xx}) > 0$ and $(\epsilon_{zz} - \epsilon_{yy}) > 0$. [18] We find that $\Delta\widetilde{\beta} = D^2\left[(\epsilon_{zz}-\epsilon_{yy})^2 - (\epsilon_{zz}-\epsilon_{xx})^2\right]$, and it can be negative ($\Delta\widetilde{\beta} < 0$) provided that ϵ_{yy} is larger than ϵ_{xx}. Furthermore, take into account the Dresselhaus system, the spin-orbit matrix $\widetilde{\beta}_S+\widetilde{\beta}_2$ yields $\Delta\widetilde{\beta} = D(\epsilon_{xx}-\epsilon_{yy})(2\beta+2\epsilon_{zz}-\epsilon_{xx}-\epsilon_{yy})$. Similar to the above case, we get $\Delta\widetilde{\beta} < 0$ when $\epsilon_{xx} < \epsilon_{yy}$. On the other hand, the system with Rashba and SIA strain split terms has the spin-orbit matrix $\widetilde{\beta}_R + \widetilde{\beta}_1$. It gives $\Delta\widetilde{\beta} = \alpha C_3(\epsilon_{xy}-\epsilon_{yx}) + \frac{1}{4}C_3^2(\epsilon_{xy}^2-\epsilon_{yx}^2)$. Theoretically, strain tensor is symmetric $\epsilon_{xy} = \epsilon_{yx}$ and $\Delta\widetilde{\beta}$ vanishes in $\widetilde{\beta}_R + \widetilde{\beta}_1$ system. However, if $\epsilon_{xy} < \epsilon_{yx}$ is experimentally exhibited, then the ISHC is expected to exceed $e/8\pi$. Consider the Dirac-Dresselhaus system whose spin-orbit matrix is given by $\widetilde{\beta}_D + \widetilde{\beta}_S$. For this system, we have $\Delta\widetilde{\beta} = 4g\beta$. This implies that if the Dresselhaus coupling β has an opposite sign to the Dirac-like coupling g, then the spin-Hall conductivity can also exceed $e/8\pi$.



FIG. 1: Variation in the spin-Hall conductivity $|\sigma_{xy}^z(0)|$ with $|N|$ for three conditions $\Delta\widetilde{\beta} > 0$, $\Delta\widetilde{\beta} = 0$, and $\Delta\widetilde{\beta} < 0$, where $Re(\lambda_</\lambda_>)$ is defined as N and $|N| < 1$.

The measurable responses caused by the spin-Hall effect are very different from those in the present idealized system, which is infinite in size and does not include impurity scattering. Measurable quantities such as spin accumulation, however, depend on boundary conditions. The conserved spin-current considered in the present paper may correspond to smooth boundaries. [28] However, the presence of impurities can drastically affect clean limit results [36–39]. In Ref. [40], it was shown that impurity scattering does not suppress the spin-Hall conductivity in the spatially random Rahsba spin-orbit coupled system. In particular, the SU(2) formulation on extrinsic mechanism of spin Hall conductivity was recently investigated in Ref. [41]. However, the effects of a finite size and impurity scattering are beyond the scope of the present paper. Hopefully, our interesting predictions of higher intrinsic ISHC would stimulate measurements in 2D semiconductor systems in the near future.

IV. ASYMMETRIC ISHC AND THE SYMMETRY OF ENERGY DISPERSION

As shown in Sec. III, the ISHC in general depends on the spin-orbit interaction, and the dependence is described by $\Delta\widetilde{\beta}$. In this section, we will show that $\Delta\widetilde{\beta}$ is related to the symmetry of energy dispersion. Therefore, we have to evaluate σ^z_{yx}. When an electric field is applied in the x direction, the spin-Hall response in the y direction is given by

$$\sigma^z_{yx}(0) = -\frac{e}{8\pi^2}\left(\det\widetilde{\beta}\right)\int_0^{2\pi} d\phi\,\frac{\sin^2\phi}{\Gamma(\phi)^2}. \tag{31}$$

The integration in Eq. (31) can also be calculated analytically as follows:

$$\sigma^z_{yx}(0) = \begin{cases} -sgn(\det\widetilde{\beta})\dfrac{e}{8\pi}\left[1 + Re\left(\dfrac{\lambda_<}{\lambda_>}\right)\right] & ;\det\widetilde{\beta} \neq 0 \\ \\ 0 & ;\det\widetilde{\beta} = 0 \end{cases} \tag{32}$$

In comparison with σ^z_{xy} [Eq. (26)], we find that σ^z_{xy} is not necessarily equal to $-\sigma^z_{yx}$ in the k-linear system. However, as mentioned in Sec. III, in the pure Rashba, pure Dresselhaus, and Rashba-Dresselhaus systems, $Re(\lambda_</\lambda_>) = 0$, and thus, $\sigma^z_{xy} = -\sigma^z_{yx}$. The asymmetric phenomenon occurs because the energy dispersion $\Gamma(\phi)$ in general does not have constrained 4-fold symmetry. Using Eq. (8), we have $\Gamma(\phi) = \Gamma(\phi + \pi)$ and

$$\begin{aligned} \Gamma(0) = \Gamma(\pi) &= \sqrt{\beta^2_{xx} + \beta^2_{yx}}, \\ \Gamma(\frac{\pi}{2}) = \Gamma(\frac{3\pi}{2}) &= \sqrt{\beta^2_{xy} + \beta^2_{yy}}. \end{aligned} \tag{33}$$

The constrained 4-fold symmetry in the energy dispersion means that the two axes k_x and k_y are equivalent in the sense that the spin splittings along k_x and k_y directions are equal. This implies $\Gamma(0) = \Gamma(\frac{\pi}{2})$, and using Eq. (28) we obtain

$$\beta^2_{xx} + \beta^2_{yx} = \beta^2_{xy} + \beta^2_{yy} \rightarrow \Delta\widetilde{\beta} = 0. \tag{34}$$

As a result, the energy dispersion with constrained 4-fold symmetry implies that the ISHC is a universal constant and the response is symmetric $\sigma^z_{xy} = -\sigma^z_{yx}$. Both the pure Rashba and pure Dresselhaus systems have circular energy dispersion, and thus, satisfy the constrained 4-fold symmetry. In the Rashba-Dresselhaus system, the energy dispersion satisfies constrained 4-fold symmetry with respect to the k_x axis and k_y axis. If the energy dispersion lacks constrained 4-fold symmetry, k_x and k_y are nonequivalent axes, thus leading to an unsymmetrical result $\sigma^z_{xy} \neq -\sigma^z_{yx}$. In this case, ISHC depends on the spin-orbit interaction ($\Delta\widetilde{\beta}$). However, it must be emphasized that the applied electric field causes a transverse spin rather than net charge current. The energy dispersion always satisfies $\Gamma(\phi) = \Gamma(\phi + \pi)$ for arbitrary ϕ since the time-reversal symmetry is not broken. This means that the electron population in $\pm k_x$ (or $\pm k_y$) is the same.

Interestingly, even if the system has constrained 4-fold symmetric energy dispersion in (k_x, k_y) axes, the asymmetric response to an applied electric field may be observed. This is because the energy dispersion may lose the constrained 4-fold symmetry in the other (k'_x, k'_y) axes. It can be shown that if $\Delta\widetilde{\beta} = 0$ the equivalent axes are at $\phi = 0, \frac{\pi}{2}, \pi, \frac{3}{2}\pi$; otherwise, they are nonequivalent axes (see Appendix B). For example, in the Rashba-Dresselhaus system, [110] and [1$\bar{1}$0] are nonequivalent axes. The corresponding spin-splittings are $(\alpha+\beta)k$ and $(\alpha - \beta)k$. We change the coordinate (k_x, k_y) to (k'_x, k'_y) such that k'_x and k'_y are parallel to [110] and [1$\bar{1}$0], respectively. The resulting effective spin-orbit matrix is

$$\begin{pmatrix} \beta_{x'x'} & \beta_{x'y'} \\ \beta_{y'x'} & \beta_{y'y'} \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}}(\alpha - \beta) & \frac{1}{\sqrt{2}}(\alpha + \beta) \\ -\frac{1}{\sqrt{2}}(\alpha - \beta) & \frac{1}{\sqrt{2}}(\alpha + \beta) \end{pmatrix}, \tag{35}$$

where x' and y' represent [110] and [$\bar{1}$10], respectively. The $\Delta\widetilde{\beta}$ corresponding to the spin-orbit matrix in Eq. (35) is $\Delta\widetilde{\beta} = -4\alpha\beta$. Using Eq. (27), we can show that $Re(\lambda_</\lambda_>) = -2\alpha\beta/[\alpha^2 + \beta^2 + |\alpha^2 - \beta^2|]$. For $\alpha^2 > \beta^2$, the ISHCs are $\sigma^z_{x'y'} = \frac{e}{8\pi}(1 + \frac{\beta}{\alpha})$ and $\sigma^z_{y'x'} = -\frac{e}{8\pi}(1 - \frac{\beta}{\alpha})$. For $\alpha^2 < \beta^2$, the ISHCs are $\sigma^z_{x'y'} = -\frac{e}{8\pi}(1 + \frac{\alpha}{\beta})$ and $\sigma^z_{y'x'} = \frac{e}{8\pi}(1 - \frac{\alpha}{\beta})$. [42] Because the determinant is independent of change of coordinate, $\det\widetilde{\beta}$ in this case is still $\alpha^2 - \beta^2$.

V. EQUILIBRIUM SPIN CURRENT AND SPIN-ORBIT MATRIX

We now turn to the discussion on equilibrium spin current in this generic k-linear spin-orbit coupled system. In Ref. [43], it was shown that even in thermodynamic equilibrium, spin current for the Rashba-Dresselhaus system does not vanish in the absence of external fields. This phenomena has arisen many discussions on the definition of spin current. [28, 44–47] The possibilities to detect the equilibrium spin currents have been studied in Refs. [45] and [48].

We calculate the equilibrium spin current by using conventional definition of spin current. In the case of the positive chemical potential ($\mu > 0$), two branches are populated. In the absence of external fields, the equilibrium spin-current is the sum of the in-plane spin currents of the two branches

$$\langle J^{\sigma_j}_i \rangle = \frac{1}{V}\sum_{n\mathbf{k}} f_{\mathbf{k}n}\langle n\mathbf{k}|\frac{1}{2}\left\{\frac{\partial H}{\partial \hbar k_i}, \frac{\hbar}{2}\sigma_j\right\}|n\mathbf{k}\rangle, \tag{36}$$

where $i, j = x, y$ and $|n\mathbf{k}\rangle$ is the eigenstate of Hamiltonian Eq. (4). From Eq. (9) and $k^+_F k^-_F = 2m\mu/\hbar^2$, a straightforward calculation yields

$$\begin{pmatrix} \langle J^{\sigma_x}_x \rangle & \langle J^{\sigma_y}_x \rangle \\ \langle J^{\sigma_x}_y \rangle & \langle J^{\sigma_y}_y \rangle \end{pmatrix} = N_F \begin{pmatrix} \beta_{yy} & -\beta_{xy} \\ -\beta_{yx} & \beta_{xx} \end{pmatrix} \det\widetilde{\beta}, \tag{37}$$

where $N_F = m^2/6\pi\hbar^4$. For specific systems, the result is in agreement with the previous results. In the pure Rashba system, where $\beta_{xy} = -\beta_{yx} = \alpha$ and $\beta_{xx} = \beta_{yy} = 0$, we have $\langle J^{\sigma_x}_y \rangle = -\langle J^{\sigma_y}_x \rangle$ and $\langle J^{\sigma_x}_x \rangle = \langle J^{\sigma_y}_y \rangle = 0$.

In the pure Dresselhaus system, where $\beta_{xx} = -\beta_{yy} = \beta$ and $\beta_{xy} = \beta_{yx} = 0$, we have $\langle J_x^{\sigma_x}\rangle = -\langle J_y^{\sigma_y}\rangle$ and $\langle J_x^{\sigma_y}\rangle = \langle J_y^{\sigma_x}\rangle = 0$. It is interesting to note that the equilibrium spin current $\langle J_i^{\sigma_j}\rangle$ is related to the inverse of spin-orbit matrix $\widetilde{\beta}^{-1}$ via $\langle J_i^{\sigma_j}\rangle = N_F(\widetilde{\beta}^{-1})_{ij}(\det\widetilde{\beta})^2$.

We find that $\det\widetilde{\beta}$ also appears in the expression of equilibrium spin current Eq. (37). However, $J_i^{\sigma_j}$ occurs in the third order of β_{ij}. In this sense, Eq. (19) fails to explain the physical meaning of $\det\widetilde{\beta}$ in this case (see Appendix). Recently, the equilibrium spin current in k-linear spin-orbit coupled systems is found to be link to the non-Abelian SU(2) gauge theory, where the Pauli spin matrix serves as a color index in the gauge field. [49] The resulting color current satisfies covariant conservation. The equilibrium spin current obtained from the covariant conserved color current in the Rashba-Dresselhaus systems is in agreement with Ref. [43]. In the following, we apply this formalism to the generic k-linear systems.

The Hamiltonian Eq. (4) can be written in terms of SU(2) gauge field $\hbar^2(\mathbf{k}+\mathbf{A})^2/2m$, where $\mathbf{A} = A_i^a\sigma_a$. We have

$$\begin{aligned} A_x^x &= \frac{m}{\hbar^2}\beta_{xx},\ A_y^x = \frac{m}{\hbar^2}\beta_{xy} \\ A_x^y &= \frac{m}{\hbar^2}\beta_{yx},\ A_y^y = \frac{m}{\hbar^2}\beta_{yy}. \end{aligned} \tag{38}$$

The equilibrium spin current denoted as J_i^a [49] is proportional to $-i\epsilon_{abc}A_i^bF_{ij}^c$, where $F_{ij}^a\sigma_a = -i[\mathbf{A}_i, \mathbf{A}_j]$ is the field strength.

The physical meaning of $\det\widetilde{\beta}$ in equilibrium spin current is now clear. The field strength in the SU(2) non-Abelian gauge field is given by [49]

$$\begin{aligned} F_{xy}^a\sigma_a &= -i[\mathbf{A}_x, \mathbf{A}_y] \\ &= -i[A_x^x\sigma_x + A_x^y\sigma_y, A_y^x\sigma_x + A_y^y\sigma_y] \\ &= -i\left(A_x^xA_y^y[\sigma_x,\sigma_y] + A_x^yA_y^x[\sigma_y,\sigma_x]\right) \\ &= 2\sigma_z\frac{m^2}{\hbar^4}\left(\det\widetilde{\beta}\right), \end{aligned} \tag{39}$$

where Eq. (38) is used. We have $F_{xy}^x = F_{xy}^y = 0$ and $F_{xy}^z = \frac{m^2}{\hbar^4}\left(\det\widetilde{\beta}\right)$. That is, $\det\widetilde{\beta}$ plays the role of color magnetic field strength F_{xy}^z.

VI. CONCLUSIONS

In conclusion, we have shown that in 2D and k-linear spin-orbit coupled systems, the properties of intrinsic spin-Hall conductivity are governed by two quantities: $\det\widetilde{\beta}$ and $\Delta\widetilde{\beta}$. The determinant of the spin-orbit matrix ($\det\widetilde{\beta}$) is a discriminant for determining whether the spin-Hall conductivity vanishes. $\det\widetilde{\beta}$ is physically expressed as the effective coupling of spin s_z and orbital motion L_z, and it explains the physical origin of the sign change of spin-Hall conductivity.

Furthermore, the dependence of spin-orbit interaction on the spin-Hall effect is related to the quantity $\Delta\widetilde{\beta}$. The quantity $\Delta\widetilde{\beta}$ reflects the symmetry of the energy dispersion. The vanishing $\Delta\widetilde{\beta}$ corresponds to the system with constrained 4-fold symmetry, and the resulting ISHC is a universal constant. However, the non-zero $\Delta\widetilde{\beta}$ gives rise to the dependence of spin-orbit interaction on ISHC. Furthermore, the sign of $\Delta\widetilde{\beta}$ changes the upper bound value of spin-Hall conductivity. For a non-vanishing spin-Hall conductivity, we show that there always exists two maximum values that the spin-Hall conductivity can achieve: $e/8\pi$ and $e/4\pi$. The conditions under which certain systems achieve a maximum value of $e/4\pi$ are also discussed.

In addition, we show that the equilibrium spin current is proportional to $(\widetilde{\beta}^{-1})_{ij}(\det\widetilde{\beta})^2$, and $\det\widetilde{\beta}$ determines the field strength of SU(2) non-Abelian gauge field in equilibrium spin current.

ACKNOWLEDGMENTS

We thank the National Science Council of Taiwan for the support under Contract No. NSC 101-2112-M-110-013-MY3.

Appendix A: Foldy-Wouthuysen transformation

A unitary transformation can be generally written as $U = e^{-iS}$, where the hermitian matrix S can be expend in order of β_{ij}, i.e. $S = S^{(1)} + S^{(2)} + S^{(3)} + \cdots$. That is, $S^{(1)}$ represents the term proportional to the order of β_{ij}, $S^{(2)}$ the order of β_{ij}^2 and so on. Follow the approach of Foldy-Wouthuysen transformation, the Hamiltonian $H = \varepsilon_{\mathbf{k}} + H_{so}$ under the unitary transformation $U = e^{-iS}$ is given by

$$\begin{aligned} H' &= e^{iS}He^{-iS} \\ &= \varepsilon_{\mathbf{k}} + H^{(1)} + H^{(2)} + H^{(3)} + o(\beta_{ij}^4), \end{aligned} \tag{A1}$$

where

$$\begin{aligned} H^{(1)} &= H_{so} + [iS^{(1)}, \varepsilon_{\mathbf{k}}] \\ H^{(2)} &= [iS^{(2)}, \varepsilon_{\mathbf{k}}] + [iS^{(1)}, H_{so}] + \frac{1}{2!}[iS^{(1)}, [iS^{(1)}, \epsilon_{\mathbf{k}}]] \end{aligned} \tag{A2}$$

and

$$\begin{aligned} H^{(3)} =& [iS^{(3)}, \epsilon_{\mathbf{k}}] + [iS^{(2)}, H_{so}] + \frac{1}{2!}[iS^{(1)}, [iS^{(2)}, \varepsilon_{\mathbf{k}}]] \\ &+ \frac{1}{2!}[iS^{(2)}, [iS^{(1)}, \varepsilon_{\mathbf{k}}]] + \frac{1}{2!}[iS^{(1)}, [iS^{(1)}, H_{so}]] \\ &+ \frac{1}{3!}[iS^{(1)}, [iS^{(1)}, [iS^{(1)}, \varepsilon_{\mathbf{k}}]]]. \end{aligned} \tag{A3}$$

Because H_{so} is an odd matrix, we have to find a matrix $S^{(1)}$ to cancel this term. Namely, we require $H^{(1)} = 0$ and

$$H_{so} + [iS^{(1)}, \varepsilon_{\mathbf{k}}] = 0. \tag{A4}$$

On the other hand, we note that H_{so} is made up of the linear momentum k_i, i.e., $H_{so} = \sigma_x(\beta_{xx}k_x + \beta_{xy}k_y) + \sigma_y(\beta_{yx}k_x + \beta_{yy}k_y)$ and $\varepsilon_{\mathbf{k}}$ is proportional to k^2, $S^{(1)}$ is obtained by the replacements $k_x \to x$ and $k_y \to y$ in H_{so}. Take into account the constant $\frac{\hbar^2}{m}$, we have

$$iS^{(1)} = \frac{im}{\hbar^2}\left\{\sigma_x(\beta_{xx}x + \beta_{xy}y) + \sigma_y(\beta_{yx}x + \beta_{yy}y)\right\}. \tag{A5}$$

Substitute Eq. (A5) into $H^{(2)}$, after a straightforward calculation, we find that the last two terms $[iS^{(1)}, H_{so}] + \frac{1}{2!}[iS^{(1)}, [iS^{(1)}, \epsilon_{\mathbf{k}}]]$ gives a diagonalized form $i\sum_{ij}\beta_{ij}^2 + \frac{2i}{\hbar}\left(\det\widetilde{\beta}\right)\sigma_z L_z$. This means that $H^{(2)}$ is already diagonalized, and thus, we can choose

$$iS^{(2)} = 0. \tag{A6}$$

Substitute Eqs. (A5) and (A6) into $H^{(3)}$ [Eq. (A3)], we obtain

$$\begin{aligned} H^{(3)} =& [iS^{(3)}, \epsilon_{\mathbf{k}}] \\ & - \frac{2m^2}{3\hbar^4}\left(\det\widetilde{\beta}\right)[\{x, L_z\}(\sigma_y\beta_{xx} - \sigma_x\beta_{yx}) \\ & + \{y, L_z\}(\sigma_y\beta_{xy} - \sigma_x\beta_{yy})]. \end{aligned} \tag{A7}$$

We find that the term in $[\cdots]$ is composed of odd matrices. Therefore, we must require $H^{(3)} = 0$. In this sense, the next diagonalized part is order of β_{ij}^4.

Appendix B: equivalent axes

In Sec. IV, we have shown that the constrained 4-fold symmetry leads to vanishing $\Delta\widetilde{\beta}$. In this appendix, we will show that when $\Delta\widetilde{\beta} = 0$, the equivalent axes are at $\phi = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}$. For equivalent axes, the locations of the axes in spherical coordinate ϕ must satisfy $\Gamma(\phi) = \Gamma(\phi + \frac{\pi}{2})$. Using Eq. (8), we have

$$\begin{aligned} \Gamma(\phi + \frac{\pi}{2})^2 =& (\beta_{xx}^2 + \beta_{yx}^2)\sin^2\phi + (\beta_{xy}^2 + \beta_{yy}^2)\cos^2\phi \\ & - (\beta_{xx}\beta_{xy} + \beta_{yx}\beta_{yy})\sin(2\phi), \end{aligned}$$

and $\Gamma(\phi) = \Gamma(\phi + \frac{\pi}{2})$ gives

$$\begin{aligned} & \left(\beta_{xx}^2 + \beta_{yx}^2 - \beta_{xy}^2 - \beta_{yy}^2\right)\cos^2\phi \\ & + \left(\beta_{xx}^2 + \beta_{yx}^2 - \beta_{xy}^2 - \beta_{yy}^2\right)\sin^2\phi \\ & + 2(\beta_{xx}\beta_{xy} + \beta_{yx}\beta_{yy})\sin(2\phi) = 0. \end{aligned} \tag{B1}$$

Substituting $\Delta\widetilde{\beta} = \beta_{xx}^2 + \beta_{yx}^2 - \beta_{xy}^2 - \beta_{yy}^2$ [Eq. (28)] into Eq. (B1), we have

$$\Delta\widetilde{\beta}\cos(2\phi) + 2(\beta_{xx}\beta_{xy} + \beta_{yx}\beta_{yy})\sin(2\phi) = 0. \tag{B2}$$

If $(\beta_{xx}\beta_{xy} + \beta_{yx}\beta_{yy})$ is non-zero, we have

$$\tan(2\phi) = -\frac{\Delta\widetilde{\beta}}{2(\beta_{xx}\beta_{xy} + \beta_{yx}\beta_{yy})}. \tag{B3}$$

For non-vanishing $\Delta\widetilde{\beta}$, the equivalent axes can be obtained by solving Eq. (B3). In particular, for vanishing $\Delta\widetilde{\beta}$, we have $\tan(2\phi) = 0$, and the locations of equivalent axes are $\phi = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}$. Because in general $\Gamma(\phi)$ satisfies $\Gamma(\pi) = \Gamma(\phi + \pi)$, the equivalent axes are $\phi = 0$ (k_x axis) and $\phi = \frac{\pi}{2}$ (k_y axis). The result also implies that, except for k_x and k_y, other axes are nonequivalent. In other words, for other arbitrary axes (k_x', k_y') that are different from (k_x, k_y), we must have $\sigma_{x'y'}^z \neq -\sigma_{y'x'}^z$. Conversely, in the pure Rashba (or pure Dresselhaus case) case, $(\beta_{xx}\beta_{xy} + \beta_{yx}\beta_{yy}) = 0$ and $\Delta\widetilde{\beta} = 0$. In this case, we have to use Eq. (B2) and we find that ϕ, the location of equivalent axes, can be arbitrary.

---

[1] G. A. Prinz, Science **282**, 1660 (1998); S. A. Wolf, D. D. Awschalom, R. A. Buhrman, J. M. Daughton, S. von Molnar, M. L. Roukes, A.Y. Chtchelkanova, and D. M. Treger, Science **294**, 1488 (2001).

[2] I. Zutic, J. Fabian, and S. D. Sarma, Rev. Mod. Phys. **76**, 323 (2004).

[3] J. E. Hirsch, Phys. Rev. Lett. **83**, 1834 (1999).

[4] S. Murakami, N. Nagaosa, and S.-C. Zhang, Science **301**, 1348 (2003); S. Murakami, N. Nagaosa, S.-C. Zhang, Phys. Rev. B **69**, 235206 (2004).

[5] J. Sinova, D. Culcer, Q. Niu, N. A. Sinitsyn, T. Jungwirth, and A. H. MacDonald, Phys. Rev. Lett. **92**, 126603 (2004).

[6] Y. K. Kato, R. C. Myers, A. C. Gossard, and D. D. Awschalom, Science **306**, 1910 (2004).

[7] N. P. Stern, S. Ghosh, G. Xiang, M. Zhu, N. Samarth and D. D. Awschalom, Phys. Rev. Lett. **97**, 126603 (2006).

[8] V. Sih, R. C. Myers, Y. K. Kato, W. H. Lau, A. C. Gossard, and D. D. Awschalom, Nature Physics **1**, 31 (2005).

[9] H. Engel, E. I. Rashba and B. I. Halperin, Phys. Rev. Lett. **95**, 166605 (2005).

[10] J. Wunderlich, et al, Phys. Rev. Lett. **94**, 047207 (2005).

[11] B. A. Bernevig and S.-C. Zhang, Phys. Rev. Lett. **95**, 016801 (2005).

[12] H. J. Chang, T.-W. Chen, J. W. Chen, W. C. Hong, W. C. Tsai, Y. F. Chen and G. Y. Guo, Phys. Rev. Lett. **98**, 136403 (2006).

[13] E. Saitoh, M. Ueda, H. Miyajima and G. Tatara, Appl. Phys. Lett. **88**, 182509 (2006); S. O. Valenzuela and M.

Tinkham, Nature (London) **442**, 176 (2006); K. Ando, Y. Kajiwara, S. Takahashi, S. Maekawa, K. Takemoto, M. Takatsu and E. Saitoh, Phys. Rev. B **78**, 014413 (2008).

[14] T. Kimura, Y. Otani, T. Sato, S. Takahashi, and S. Maekawa, Phys. Rev. Lett. **98**, 156601 (2007).

[15] G. Y. Guo, S. Murakami, T.-W. Chen and Nagaosa, Phys. Rev. Lett. **100**, 096401 (2008).

[16] A. Bohm, A. Mostafazadeh, H. Koizumi, Q. Niu and J. Zwanziger, *The Geometric Phase in Quantum Systems* (Springer, Berlin, 2003).

[17] T. Seki, Y. Hasegawa, S. Mitani, S. Takahashi, H. Imamura, S. Maekawa, J. Nitta and K. Takahashi, Nature Mater. **7**, 125 (2008).

[18] B. A. Bernevig and S.-C. Zhang, Phys. Rev. B **72**, 115204 (2005); Y. Kato, R. C. Myers, A. C. Gossard, D. D. Awschalom, Nature (London) **427**, 50 (2004); Y. Kato, R. C. Myers, A. C. Gossard, D. D. Awschalom, Phys. Rev. Lett. **93**, 176601 (2004).

[19] S.-Q. Shen, Phys. Rev. B **70**, 081311(R) (2004); N. A. Sinitsyn, E. M. Hankiewicz, W. Teizer, and J. Sinova, Phys. Rev. B **70**, 081312(R) (2004); M.-C. Chang, Phys. Rev. B **71**, 085315 (2005); T.-W. Chen, C.-M. Huang and G. Y. Guo, Phys. Rev. B **73**, 235309 (2006).

[20] J. Nitta, T. Akazaki, H. Takayanagi and T. Enoki, Phys. Rev. Lett. **78**, 1335 (1997).

[21] E. I. Rashba, Sov. Phys. Solid State **2**, 1224 (1960); Y. A. Bychkov and E. I. Rashba, J. Phys. C **17**, 6039 (1984).

[22] G. Dresselhaus, Phys. Rev. **100**, 580 (1955).

[23] J. W. McClure, Phys. Rev. **104**, 666 (1956); D. P. DiVincenzo and E. J. Mele, Phys. Rev. B **29**, 1685 (1984); V. P. Gusynin and S. G. Sharapov, **95**, 146801 (2005); 19.K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A. Firsov, Science **306**, 666 (2004); K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, Nature (London) **438**, 197 (2005); Y. Zhang, J. P. Small, M. E. S. Amori, and P. Kim, Phys. Rev. Lett. **94**, 176803 (2005); Y.-W. Tan, H. L. Stormer, and P. Kim, Nature (London) **438**, 201 (2005); Y. Zhang, Y.-W. Tan, H. L. Stormer, and P. Kim, **438**, 201 (2005).

[24] N. A. Sinitsyn, A. H. MacDonald, T. Jungwirth, V. K. Dugaev and J. Sinova, Phys. Rev. B **75**, 045315 (2007).

[25] G. E. Pikus and A. N. Titkov, *Optical Orientation*(North-Holland, Amsterdam. 1984), p. 73.

[26] In constructing the spin-orbit matrix $\widetilde{\beta}_2$, we have neglected the term $Dk_z(\epsilon_{yy} - \epsilon_{xx})$ which exists when $\epsilon_{xx} \neq \epsilon_{yy}$. However, since k_z is not influenced by the in-plane electric field, $\langle k_z \rangle = 0$, we can approximately neglect this term. (see also [18])

[27] G. D. Mahan, *Many-Particle Physics* (Kluwer Academic/Plenum publisher, New York, 2000).

[28] D. Culcer, J. Sinova, N. A. Sinitsyn, T. Jungwirth and A. H. MacDonald and Q. Niu, Phys. Rev. Lett. **93**, 046602 (2004); J. Shi, P. Zhang, D. Xiao and Q. Niu, Phys. Rev. Lett. **96**, 076604 (2006).

[29] B. A. Bernevig, Phys. Rev. B **71**, 073201 (2005).

[30] M. Valin-Rodriguez, Phys. Rev. Lett. **107**, 266801 (2011).

[31] L. L. Foldy and S. A. Wouthuysen, Phys. Rev. **78**, 29 (1950); T.-W. Chen and D.-W. Chiou, Phys. Rev. A. **82**, 012115 (2010).

[32] G. Arfken and H. J. Weber, *Mathematical Methods for Physicists* (Academic Press, 1995).

[33] E. I. Rashba, Physica E: Low-dimensional Systems and Nanostructures **34**, 31 (2006); Hyun C. Lee and S.-R. Eric Yang, Phys. Rev. B **72**, 245338 (2005).

[34] B. Jusserand, D. Richards, G. Allan, C. Priester and B. Etienne, Phys. Rev. B **51**, 4707 (1995); W. Knap, C. Skierbiszewski, A. Zduniak, E. Litwin-Staszewska, D. Bertho, F. Kobbi, J. L. Robert, G. E. Pikus, F. G. Pikus, S. V. Iordanskii, V. Mosser, K. Zekentes, and Yu. B. Lyanda-Geller, Phys. Rev. B **53**, 3912 (1996); J. B. Miller, D. M. Zumbuhl, C. M. Marcus, Y. B. Lyanda-Geller, D. Goldhaber-Gordon, K. Campman, and A. C. Gossard, Phys. Rev. Lett. **90**, 076807 (2003).

[35] T.-W. Chen and G. Y. Guo, Phys. Rev. B **79**, 125301 (2009).

[36] J. Sinova, S. Murakami, S.-Q. Shen and M.-S. Choi, Solid State Commun. **138**, 214 (2006); J. I. Inoue, G. E. W. Bauer and L. W. Molenkamp, Phys. Rev. B **70**, 041303(R) (2004); E. G. Mishchenko, A. V. Shytov and B. I. Halperin, Phys. Rev. Lett. **93**, 226602 (2004); O. Chalaev and D. Loss, Phys. Rev. B **71**, 245318 (2005); O. V. Dimitrova, Phys. Rev. B **71**, 245327 (2005).

[37] J. Schliemann and D. Loss, Phys. Rev. B **69**, 165315 (2004); A. G. Mal'shukov and K. A. Chao, Phys. Rev. B**71**, 121308 (R) (2005); A. Khaetskii, Phys. Rev. B **73**, 115323 (2006); N. Sugimoto, S. Onoda. S. Murakami and N. Nagaosa, Phys. Rev. B **73**, 113305 (2006).

[38] A. A. Burkov, Alvaro S. Nunez and A. H. MacDonald, Phys. Rev. B **70**, 155308 (2004).

[39] K. Nomura, J. Sinova, T. Jungwirth, Q. Niu, and A. H. MacDonald, Phys. Rev. B **71**, 041304 (2005); B. K. Nikolic, L. P. Zarbo, and S. Souma, Phys. Rev. B **72**, 075361 (2005); L. Sheng, D. N. Sheng, C. S. Ting, Phys. Rev. Lett. **94**, 016602 (2005); E. M. Hankiewicz, L. W. Molenkamp, T. Jungwirth, and J. Sinova, Phys. Rev. B **70**, 241301(R) (2004).

[40] V. K. Duagev, M. Inglot, E. Ya. Sherman, and J. Barna$\acute{s}$, Phys. Rev. B **82**, 121310(R) (2010).

[41] R. Raimondi, P. Schwab, C. Gorini, and G. Vignale, Ann. Phys. (Berlin) 524, 153 (2012) [arXiv:1110.5279].

[42] T.-W. Chen, H.-C. Hsu and G. Y. Guo, Phys. Rev. B **80**, 165302 (2009).

[43] E. I. Rashba, Phys. Rev. B **68**, 241315(R) (2003).

[44] E. I. Rashba, Phys. Rev. B **70**, 161201(R) (2004); A. A. Burkov, A. S. N$\acute{u}\tilde{n}$ez and A. H. MacDonald, Phys. Rev. B **70**, 155308 (2004); E. B. Sonin, Phys. Rev. B **76**, 033306 (2007).

[45] Q.-f. Sun, X. C. Xie and J. Wang, Phys. Rev. Lett. **98**, 196801 (2007); Phys. Rev. B **77**, 035327 (2008).

[46] V. A. Sablikov, A. A. Sukhanov and Y. Ya. Tkach, Phys. Rev. B **78**, 153302 (2008).

[47] H.-J. Drouhin, G. Fishman and J.-E. Wegrowe, Phys. Rev. B **83**, 113307 (2011).

[48] E. B. Sonin, Phys. Rev. Lett. **99**, 266602 (2007).

[49] I. V. Tokatly, Phys. Rev. Lett. **101**, 106601 (2008).